

Intercluster Correlation in Seismicity

Mega et al. [1] proposed to use the “diffusion entropy” (DE) method to demonstrate that the distribution of time intervals between a large earthquake (the mainshock of a given seismic sequence) and the next one does not obey Poisson statistics. We have re-analyzed the same catalog of seismic events in California and performed synthetic tests which show that the DE is unable to detect correlations between clusters, thus negating the claimed possibility of detecting an intercluster correlation. We have generated synthetic earthquake catalogs with a Poisson distribution of mainshock times, with aftershocks triggered in direct lineage by each mainshock following Omori’s power-law decay of seismic rate $n(t) \propto 1/(t - t_i)^p$ (with $p = 1$, and where t_i is the mainshock time) and with a power law distribution of aftershock cluster sizes with exponent $b/\alpha = 1.25$ (justified from the interplay between the power-law distribution of energies with exponent $b = 1$ (Gutenberg-Richter law) and the power-law increase of aftershock productivity with the mainshock energy of exponent $\alpha = 0.8$ [2]). Fig. 1 (crosses) shows that our synthetic DE without intercluster correlation is essentially undistinguishable from the real data. Thus, the conclusion of Mega et al. [1] that the clusters of seismicity are correlated is not warranted by their analysis. However, the *fact* that intercluster correlation exists is known at least since Kagan and Jackson [3] and can be observed straightforwardly, as shown in Fig. 2.

We have used the ETAS model to generate realistic synthetic catalogs. The ETAS model is a simple branching model of seismicity [4] which contains the Gutenberg-Richter, the Omori law, the productivity law, cascades of multiple triggering between earthquakes and a poissonian seismicity background. In the ETAS model, any earthquake may trigger other earthquakes, without arbitrary distinction between foreshocks, aftershocks and mainshocks. The ETAS model reproduces many properties of seismicity, including realistic foreshock sequences [5]. It is widely used to model and predict the spatio-temporal distribution of seismicity (e.g., [4] and ref. 8 of [1]). As shown in Figs. 1 and 2 (circles), the ETAS model recovers (i) Mega et al.’s observation of DE $S(t) = A + \delta \ln t$ with $\delta = 0.94$, (ii) the power-law pdf of the time intervals $\tau^{[m]}$ between two successive mainshocks (which arises because many mainshocks are also aftershocks of other mainshocks) and (iii) a correlation between $\tau_i^{[m]}$ ’s (which is absent in Mega et al.’s LR model). Thus, their LR model, which introduces correlation between clusters of seismicity, is insufficient to account for the correlation observed in the data.

We believe that these discrepancies with Mega et al.’s conclusions stem from their incorrect use of the pdf of intracuster interevent times $t_{j+1} - t_j$. Omori’s law is the pdf of the times between a mainshock and its aftershocks and describes a non-stationary process such




FIG. 1: The DE $S(t)$ defined in [1] calculated for a Poisson process (squares), the California catalog used in [1] (diamonds), the GP model used in [1] and the ETAS model (circles) with parameters: branching ratio $n = 0.93$, $b = 1$, Omori exponent $p = 1.3$ for first-generation triggering and $\alpha = 0.8$.




FIG. 2: (a) probability density function (pdf) of the time intervals $\tau^{[m]}$ between two successive large ($M \geq 5$) earthquakes in the Southern California catalog (circles) and in the ETAS model (crosses). (b) recurrence plot $\ln \tau_{i+1}^{[m]}$ vs $\ln \tau_i^{[m]}$. The straight line is the linear fit to the data (circles). The correlation is $r = 0.43$ (significance $> 99.9\%$) for the data and $r = 0.21$ (significance $> 99\%$) for ETAS.

that the interevent times increase as time increases since the mainshock: for $p = 1$ for instance, $t_{j+1} - t_j \sim t_j$. Therefore, the pdf of interevent times is not equivalent to Omori’s law and it loses most of the information on correlations between aftershocks. This work is supported by NSF-EAR02-30429 and by the Southern California Earthquake Center (SCEC).

-
- [1] M.S. Mega et al., Phys. Rev. Lett. 90, 10.1103, 188501 (2003).
 - [2] A. Helmstetter, Phys. Rev. Lett. XX, XX (2003) (physics/0210056).
 - [3] Y.Y. Kagan and D.D. Jackson, Bull. Seism. Soc. Am. 89, 1147 (1999).
 - [4] Y.Y. Kagan and L. Knopoff, J. Geophys. Res. 86, 2853 (1981); Y. Ogata, J. Am. stat. Assoc. 83, 9 (1988).
 - [5] A. Helmstetter and D. Sornette, in press in J. Geophys. Res. (physics/0210130).

A. Helmstetter¹ and D. Sornette^{1,2}, ¹ Institute of Geo-

physics and Planetary Physics, UCLA ² Department of Earth and Space Sciences, University of California, Los Angeles, California 90095-1567 and LPMC, CNRS UMR 6622 and Université de Nice, 06108 Nice, France